A prospective comparative assessment of surgical outcomes following laparoscopic and conventional appendectomy

[1]Dr Adil Shah Roghani, Reema Naveed

Kabir Medical College, Peshawar.

Abstract

Background: Acute appendicitis is commonly treated by appendectomy, performed using either a laparoscopic or an open approach. The choice of technique may influence postoperative pain, wound-related complications, recovery time, length of hospital stay, and return to normal activities. Objective: To compare postoperative outcomes among patients undergoing laparoscopic appendectomy and open appendectomy. Methods: This prospective comparative observational study should enroll eligible patients undergoing appendectomy at the study site during a predefined recruitment period. Baseline characteristics, operative details, postoperative pain, complications, length of stay, readmission, and recovery milestones should be recorded using a standardized case-report form. Outcomes should be analyzed according to the prespecified statistical plan. Results: Results should be inserted after the study dataset has been collected, cleaned, and analyzed. Report the number of participants in each group, effect estimates, confidence intervals, and exact p-values where appropriate. Conclusion: The comparative conclusion must be based on the observed data and adjusted for relevant clinical differences between groups.

Keywords: appendicitis; laparoscopic appendectomy; open appendectomy; postoperative pain; surgical site infection; length of hospital stay.

[1]Alam Javed, [2]IKram Khan, [3]Dr Adil Shah Roghani, [2]Reema Naveed

Kabir medical college, Peshawar.

Submitted: June 5, 2025 Revised: January 20, 2026

Accepted: March 6, 2026

Introduction

Acute appendicitis is one of the most common causes of acute abdominal pain requiring emergency surgical intervention worldwide. Appendectomy remains the standard treatment for most patients with acute appendicitis, although the choice of surgical approach depends on disease severity, patient characteristics, surgeon expertise, and available resources. Traditionally, open appendectomy has been performed through an abdominal incision that provides direct access to the appendix. However, laparoscopic appendectomy has become increasingly popular because of advances in minimally invasive surgical techniques and improvements in perioperative care. Both approaches are widely used, and their comparative effectiveness continues to be evaluated in terms of postoperative recovery,

complications, and healthcare utilization 1. Laparoscopic appendectomy involves the insertion of a camera and specialized surgical instruments through small abdominal incisions. This technique enables visualization of the abdominal cavity and removal of the inflamed appendix while minimizing the size of the surgical wounds. Potential advantages include reduced postoperative pain, improved cosmetic outcomes, earlier mobilization, and shorter hospitalization. Furthermore, laparoscopy allows surgeons to inspect other intra-abdominal structures when the diagnosis is uncertain or additional pathology is suspected. Nevertheless, laparoscopic surgery requires specialized equipment, appropriate training, and experience. Operative duration, equipment availability, and the risk of complications associated with trocar insertion or pneumoperitoneum may also influence its suitability in particular clinical circumstances 2.

Open appendectomy remains an effective and established surgical procedure, particularly in healthcare settings where laparoscopic equipment or trained personnel are not consistently available. It may also be appropriate in selected patients with extensive intraabdominal inflammation, complex anatomical findings, or circumstances requiring conversion from laparoscopic surgery. Although the open approach provides direct access to the operative site, it generally involves a larger incision that may contribute to postoperative discomfort, wound complications, and delayed return to normal activities. The relative advantages of each technique should therefore be assessed according to measurable clinical outcomes rather than assumptions about the superiority of either procedure 3.

Postoperative outcomes are important indicators of surgical quality and patient recovery. Commonly evaluated parameters include postoperative pain, analgesic requirements, surgical-site infection, intra-abdominal abscess, operative duration, length of hospital stay, readmission, and time required to resume normal daily activities. Surgical-site infections can increase treatment costs, prolong hospitalization, and negatively affect patient satisfaction. The risk of postoperative complications may be influenced by age, obesity, diabetes mellitus, disease severity, perforation, operative technique, antibiotic administration, and adherence to infection-prevention protocols. Consequently, comparative studies should consider these variables when interpreting differences between laparoscopic and open appendectomy 4,54,5. Previous comparative investigations and systematic reviews have examined the outcomes associated with laparoscopic and open appendectomy. Evidence has frequently suggested benefits of the laparoscopic approach for selected recovery outcomes, including wound-related complications and duration of hospitalization. However, the magnitude of these benefits may vary according to patient selection, institutional resources, surgical expertise, and the proportion of patients presenting with complicated appendicitis. Differences in study design, outcome definitions, follow-up duration, and perioperative management can also contribute to inconsistent findings. Therefore, further prospective evaluations in specific clinical settings may provide useful information for

improving surgical decision-making and postoperative care 6.

A prospective observational study offers an opportunity to collect standardized clinical information before, during, and after surgery. By recording baseline characteristics, operative findings, and postoperative events systematically, researchers can compare outcomes between patients undergoing laparoscopic and open appendectomy under routine clinical conditions. Appropriate statistical analysis can help identify differences in outcomes while accounting for relevant confounding variables. Nevertheless, because surgical allocation is not randomized in an observational study, the findings should be interpreted cautiously, and associations should not automatically be considered causal 7. The present study aims to compare postoperative outcomes following laparoscopic versus open appendectomy, focusing on postoperative pain, surgical complications, length of hospital stay, and recovery. The findings may help inform local surgical practice, support evidence-based treatment decisions, and identify opportunities to improve postoperative management among patients undergoing appendectomy.

Introduction

Acute appendicitis is one of the most common causes of acute abdominal pain requiring emergency surgical intervention. It occurs when the appendix becomes inflamed, commonly due to luminal obstruction, which may lead to swelling, infection, ischemia, and, in severe cases, perforation. Early diagnosis and timely surgical treatment are essential to reduce complications and improve patient outcomes. Appendectomy remains the standard treatment for most patients with acute appendicitis, and it can be performed using either the conventional open technique or a minimally invasive laparoscopic approach \(1\).

Open appendectomy has been used for decades and remains an effective treatment option, particularly in settings where laparoscopic equipment or trained personnel are unavailable. The procedure generally involves an incision in the right lower abdominal region to access and remove the appendix. Although it is well established and can be performed safely in many clinical situations, open surgery may be associated with greater wound discomfort, a larger surgical incision, and a longer recovery period in some patients. These considerations have encouraged the adoption of laparoscopic appendectomy as an alternative surgical approach.


Laparoscopic appendectomy is performed through small abdominal incisions using a camera and specialized instruments. This technique allows the surgeon to visualize the abdominal cavity and remove the inflamed appendix while minimizing the size of the surgical wounds. Potential advantages include reduced postoperative pain, fewer wound infections, improved cosmetic outcomes, and earlier return to normal activities. However, the procedure may require specialized equipment, greater technical expertise, and additional operating-room resources. The benefits may also vary according to the severity of appendicitis, patient characteristics, and the experience of the surgical team \(2,3\).

Comparisons between laparoscopic and open appendectomy have demonstrated differences in

several postoperative outcomes, although the magnitude of these differences may vary across studies. Important measures include postoperative pain, surgical-site infection, duration of hospital stay, operative time, postoperative complications, and time required to resume routine activities. Laparoscopic surgery is frequently associated with advantages in wound-related outcomes and recovery, but the choice of technique should also consider the patient's clinical condition, available resources, and the surgeon's expertise \(2,3\). International guidelines support the use of laparoscopic appendectomy when appropriate equipment and surgical expertise are available, while recognizing that open appendectomy remains a suitable option in selected circumstances \(4\).

A prospective comparison of the two techniques can provide useful evidence regarding their performance in a defined clinical setting. By assessing postoperative outcomes systematically, researchers can identify differences in recovery, complications, and healthcare utilization. Consistent outcome definitions and appropriate follow-up are important to ensure that the findings are reliable and clinically meaningful. Study reporting should also clearly describe participant selection, surgical procedures, outcome measures, and follow-up methods \(5\).

Therefore, this study aims to compare laparoscopic and open appendectomy in terms of postoperative outcomes. The comparison focuses on postoperative pain, wound complications, length of hospital stay, operative duration, and recovery. The findings may help inform surgical decision-making and support the selection of an appropriate approach according to patient needs, clinical circumstances, and available resources. **Illustrations**






Figure 1. Conceptual comparison of laparoscopic and open appendectomy approaches. Schematic illustration; not to scale.

Figure 2. Key postoperative outcomes for prospective comparison. Outcomes should be measured using predefined criteria.

Aim and Objectives Aim: To compare postoperative outcomes following laparoscopic versus open appendectomy.

- To compare postoperative pain and analgesic requirements between groups.
- To compare surgical-site infection and other postoperative complications.
- To compare length of hospital stay and time to return to usual activities.
- To document readmission and reintervention within the predefined follow-up period.

Methodology Study design and setting

A prospective comparative observational study should be conducted in the surgical department of [hospital/institution] over [start date] to [end date]. The protocol should receive approval from the relevant institutional ethics committee before recruitment.

Study population

Eligible participants are patients undergoing appendectomy for suspected acute appendicitis during the recruitment period. Inclusion and exclusion criteria should be defined before enrollment and applied consistently to both groups. **Inclusion criteria**

- Patients undergoing laparoscopic or open appendectomy.
- Patients for whom informed consent and follow-up data can be obtained.
- Patients meeting the study protocol's predefined age and clinical eligibility criteria.

Exclusion criteria

- Patients requiring a different primary operation.
- Patients with missing essential baseline or outcome data, according to the prespecified analysis plan.
- Patients declining participation or unable to complete the required follow-up, where applicable.

Data collection and outcomes Record age, sex, body mass index where available, comorbidities, symptom duration, clinical and imaging findings, disease severity, perforation status, antibiotic use, operative duration, conversion to open surgery, and surgeon-related variables where feasible. Define postoperative pain measurement time points and scale, surgical-site infection criteria, intraabdominal abscess, length of stay, readmission, reintervention, and return-to-activity criteria before data collection.

Statistical analysis

Summarize continuous variables using mean and standard deviation or median and interquartile range, as appropriate; summarize categorical variables using counts and percentages. Use suitable comparative tests based on distribution and cell counts. Report effect estimates with 95% confidence intervals. Multivariable regression may be used to adjust for clinically important confounders. Specify the software, significance threshold, missing-data approach, and analysis population in the final manuscript. **Ethical considerations** Obtain institutional ethics approval and informed consent where required. Protect participant confidentiality, store data securely, and report the study in accordance with relevant observational-study reporting guidance.

Results

Insert results only after analysis of the actual study dataset. The tables below are structured templates and intentionally contain no fabricated values.

		)	
Age, years (mean ± SD)			
Sex, n (%)			
BMI, kg/m² (mean ± SD)			
Comorbidities, n (%)			
Symptom duration, hours			
Complicated/ perforated appendicitis, n (%)			
Drain placement, n (%)			
Perioperative antibiotic regimen			

Table 1. Baseline demographic and clinical characteristics

Variable	Laparoscopic (n=)	Open (n=	pvalue

Outcome	Laparoscopic (n=)	Open (n=)	Effect estimate (95% CI) / pvalue
Postoperative pain score at defined time			
Analgesic requirement, n (%)			

Interpretation: Complete this paragraph after statistical analysis, reporting the direction and magnitude of group differences without overstating observational associations.

Table 3. Postoperative outcomes

Interpretation: Complete this paragraph after statistical analysis, reporting the direction and magnitude of group differences without overstating observational associations. **Table 2. Operative characteristics**

Variable	Laparoscopic (n=)	Open (n=)	pvalue
Operative time, minutes			
Intraoperative complications, n (%)			
Conversion to open surgery, n (%)		Not applicable	
Length of stay, days			
Return to usual activity, days			
Surgicalsite infection, n (%)			
Intraabdominal abscess, n (%)			

Interpretation: Complete this paragraph after statistical analysis, reporting the direction and magnitude of group differences without overstating observational associations. **Table 4. Adjusted analysis of postoperative complications**

Predictor	Adjusted effect estimate	95 % CI	pvalue
Surgical approach (laparoscopic vs open)			
Age			
Comorbidity status			
Perforated/ complicated appendicitis			
Operative duration			

Interpretation: Complete this paragraph after statistical analysis, reporting the direction and magnitude of group differences without overstating observational associations.

Discussion

The present study compares postoperative outcomes following laparoscopic versus open appendectomy, focusing on postoperative pain,

surgical-site infection, postoperative complications, length of hospital stay, and recovery. The comparison is clinically important because both procedures are widely used for acute appendicitis, and the choice of surgical approach may influence patient recovery and healthcare utilization. The interpretation of the findings should be based on the actual study results, considering patient characteristics, disease severity, operative factors, and postoperative management. Laparoscopic appendectomy has gained acceptance because minimally invasive surgery may offer advantages in postoperative recovery. Smaller incisions may reduce wound trauma, postoperative discomfort, and the risk of superficial surgical-site infection. In addition, laparoscopic visualization allows surgeons to inspect the abdominal cavity and identify associated pathology when necessary. Jaschinski et al. 1, in a Cochrane systematic review, compared laparoscopic and open surgery for suspected appendicitis and evaluated outcomes including wound infection, intra-abdominal abscess, pain, and length of hospitalization. These findings provide an important evidence base for evaluating the potential benefits and limitations of both surgical approaches. Postoperative pain is another important measure of surgical effectiveness. Patients undergoing laparoscopic appendectomy may experience less incisional pain because the procedure generally requires smaller abdominal incisions. Reduced pain may facilitate early mobilization and improve patient comfort during the initial postoperative period. However, pain outcomes can also depend on analgesic protocols, operative duration, individual pain sensitivity, and disease severity. Di Saverio et al. 2 emphasized the importance of evidence-based management of acute appendicitis, including appropriate selection of surgical treatment according to clinical circumstances. Consequently, pain scores should be recorded using a standardized scale at predefined postoperative intervals to ensure reliable comparisons between groups. Surgical-site infection and other postoperative complications are essential indicators of surgical quality. Open appendectomy involves a larger incision, which may increase exposure of the wound to contamination and contribute to wound-related morbidity. Nevertheless, laparoscopic appendectomy is not free from complications, including intra-abdominal abscess, particularly in patients with complicated or perforated appendicitis. The Centers for Disease Control and Prevention guidelines highlight the importance of appropriate perioperative antimicrobial prophylaxis and standardized infection-prevention practices 3. Differences in antibiotic administration, operative technique, and postoperative wound care should therefore be considered when interpreting infection rates. Length of hospital stay and time to return to normal activities are also relevant outcomes because they influence patient satisfaction, hospital resource utilization, and overall treatment costs. A shorter hospital stay may reflect reduced postoperative pain, fewer wound complications, and faster functional recovery. However, discharge decisions may vary according to institutional protocols, social circumstances, and the availability of outpatient follow-up. The World Health Organization 4

recommends evidence-based measures to prevent surgical-site infections, supporting the importance of standardized perioperative care when evaluating postoperative outcomes. The interpretation of comparative outcomes should also account for methodological limitations. In a prospective observational study, the surgical approach may be influenced by disease complexity, surgeon preference, equipment availability, and patient characteristics. These differences may introduce confounding and affect the observed associations. Bhangu et al. 5 discussed the complexities of diagnosing and managing acute appendicitis, reinforcing the importance of considering clinical presentation and disease severity in treatment decisions. Appropriate multivariable analysis can help account for measured confounders, although residual confounding may remain. Overall, the comparison of laparoscopic and open appendectomy should integrate postoperative complications, pain, hospitalization, and recovery rather than rely on a single outcome. The final interpretation must be aligned with the observed results, and conclusions should not imply superiority unless supported by the study data. Observational research should also be reported transparently in accordance with established reporting recommendations 6.

Conclusion

The relative postoperative outcomes of laparoscopic and open appendectomy should be concluded from the study's analyzed data. The final conclusion should state the principal observed differences, their uncertainty, and the clinical context, while acknowledging the limitations of a prospective observational design.

References

1. Bhangu, A., Søreide, K., Di Saverio, S., Assarsson, J. H., & Drake, F. T. (2015). Acute appendicitis: Modern understanding of pathogenesis, diagnosis, and management. The Lancet, 386(10000), 1278–1287. doi.org
2. Di Saverio, S., Podda, M., De Simone, B., Ceresoli, M., Augustin, G., Gori, A., et al. (2020). Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. World Journal of Emergency Surgery, 15, 27. doi.org
3. Jaschinski, T., Mosch, C. G., Eikermann, M., Neugebauer, E. A. M., & Sauerland, S. (2018). Laparoscopic versus open surgery for suspected appendicitis. Cochrane Database of Systematic Reviews, 2018(1), CD001546. doi.org
4. Berríos-Torres, S. I., Umscheid, C. A., Bratzler, D. W., Leas, B., Stone, E. C., Kelz, R. R., et al. (2017). Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017. JAMA Surgery, 152(8), 784–791. doi.org
5. World Health Organization. (2018). Global guidelines for the prevention of surgical site infection (2nd ed.). World Health Organization. who.int
6. Jaschinski, T., et al. (2018). Laparoscopic versus open surgery for suspected appendicitis. Cochrane Database of Systematic Reviews, 2018(1), CD001546. doi.org

7. von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., & Vandenbroucke, J. P. (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. PLoS Medicine, 4(10), e296. doi.org.
8. Bhangu, A., Søreide, K., Di Saverio, S., Assarsson, J. H., & Drake, F. T. (2015). Acute appendicitis: Modern understanding of pathogenesis, diagnosis, and management. The Lancet, 386(10000), 1278–1287. doi.org
9. Jaschinski, T., Mosch, C. G., Eikermann, M., Neugebauer, E. A. M., & Sauerland, S. (2018). Laparoscopic versus open surgery for suspected appendicitis. Cochrane Database of Systematic Reviews, 2018(11), CD001546. doi.org
10. Jaschinski, T., Mosch, C., Eikermann, M., & Neugebauer, E. A. M. (2015). Laparoscopic versus open appendectomy in patients with suspected appendicitis: A systematic review of metaanalyses of randomized controlled trials. BMC Gastroenterology, 15, 48. doi.org).
11. Di Saverio, S., Podda, M., De Simone, B., Ceresoli, M., Augustin, G., Gori, A., et al. (2020). Diagnosis and treatment of acute appendicitis: 2020 update of the WSES Jerusalem guidelines. World Journal of Emergency Surgery, 15, 27. doi.org
12. Berríos-Torres, S. I., Umscheid, C. A., Bratzler, D. W., Leas, B., Stone, E. C., Kelz, R. R., et al. (2017). Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017. JAMA Surgery, 152(8), 784–791. doi.org